

Survey for Supernovae in Massive High-Redshift Clusters

Keren Sharon^{*}, Avishay Gal-Yam^{†,**}, Dan Maoz^{*}, Megan Donahue[‡], Harald Ebeling[§], Richard S. Ellis[†], Alex V. Filippenko[¶], Ryan Foley[¶], Wendy L. Freedman^{||}, Robert P. Kirshner^{††}, Jean-Paul Kneib^{‡‡}, Thomas Matheson^{§§}, John S. Mulchaey^{||}, Vicki L. Sarajedini^{¶¶} and Mark Voit[‡]

^{*}*School of Physics and Astronomy, Tel Aviv University*

[†]*Division of Physics, Mathematics and Astronomy, California Institute of Technology*

^{**}*Hubble Fellow*

[‡]*Michigan State University*

[§]*University of Hawaii*

[¶]*University of California - Berkeley*

^{||}*Carnegie Institution of Washington*

^{††}*Harvard University*

^{‡‡}*Observatoire de Marseille*

^{§§}*National Optical Astronomy Observatories, AURA*

^{¶¶}*University of Florida*

Abstract. We describe our ongoing program designed to measure the SN-Ia rate in a sample of massive $z=0.5-0.9$ galaxy clusters. The SN-Ia rate is a poorly known observable, especially at high z , and in cluster environments. The SN rate and its redshift dependence can serve as powerful discriminants for a number of key issues in astrophysics and cosmology. Our observations will put clear constraints on the characteristic SN-Ia “delay time”, the typical time between the formation of a stellar population and the explosion of some of its members as SNe-Ia. Such constraints can exclude entire categories of SN-Ia progenitor models, since different models predict different delays. These data will also help to resolve the question of the dominant source of the high metallicity in the intracluster medium (ICM) - SNe-Ia, or core-collapse SNe from an early stellar population with a top-heavy IMF, perhaps those population III stars responsible for the early re-ionization of the Universe. Since clusters are excellent laboratories for studying enrichment (they generally have a simple star-formation history, and matter cannot leave their deep potentials), the results will be relevant for understanding metal enrichment in general, and the possible role of first generation stars in early Universal enrichment. Observations obtained so far during cycles 14 and 15 yield many SNe in our cluster fields, but our follow-up campaign reveals most are not in cluster galaxies.

Keywords: Supernovae

PRELIMINARY RESULTS

We were allocated 30 orbits of *HST* time during cycles 14 and 15¹, to re-image 15 X-ray luminous clusters, which were already imaged by ACS in the past (e.g., Fig. 1). We compare our new images with previous data, and promptly search for transients using image subtraction (Fig. 2).

¹ see <http://www.astro.caltech.edu/~avishay/hst.html> for additional details

TABLE 1. Candidate SNe from our *HST* search.

SN	Magnitude	Redshift	ID
M0257_1	25.3	?	
M0257_2	27.4	?	Hostless SN
M0257_3	26.1	0.73	Background SN
M0257_4	26.4	?	
M0257_5	25.7	?	
M0647_1	25.8	?	
M0647_2	26.8	?	
M0717_1	25.8	0.55	Cluster, Non-Ia SN
M0911_1	24.3	?	
M0911_2	24.9	?	
M0911_3	24.0	?	
M1149_1	23.9	?	
M1149_2	24.6	?	
M1149_3	24.7	?	
M2129_1	24.9	0.87	Background SN
M2129_2	24.4	?	Hostless SN
M2214_1	26.2	0.58	Background SN
M2214_2		?	
M2214_3	24.1	?	
M2214_4		?	
C1226_1	25.8	?	
C1226_2	24.0	0.90	Cluster SN Ia
M1054_1	24.8	?	
M1054_1	23.4	0.83	Cluster SN Ia
M0451_1	25.5	0.16	Foreground SN
C0152_1	25.2	1.27	AGN
C0152_2	>24	?	

Follow-up spectroscopy (e.g., Fig. 3) is used to determine host redshifts for faint SNe (typical events have $I > 25$ mag) and thus reject foreground or background events and AGN. Color imaging is used to further classify SNe based on their colors and luminosities. Partial follow-up has already been secured, and additional Keck follow-up has been allocated. Table 1 lists the SN candidates that were found in Cycle 14. For each event we give an identification (if possible). Two cluster SNe Ia have been secured so far, and additional 2-6 (hostless, or in galaxies which appear to be cluster members based on color and *HST* morphology) should also contribute to the cluster rate. All other events appear to reside in galaxies which are most likely interlopers and AGN.

ACKNOWLEDGMENTS

A.G. acknowledges support by NASA through Hubble Fellowship grant #HST-HF-01158.01-A awarded by STScI, which is operated by AURA, Inc., for NASA, under contract NAS 5-26555. A.G. further acknowledges the hospitality of the community of Cefalu and the efforts of the organizers of the 2006 Cefalu international astronomy conference, during which this work has come to fruition.




FIGURE 1. Example of an ACS cluster image from our ongoing program, with zoom-ins on two of the SNe we have found.




FIGURE 2. Examples of SNe and SN candidates found in our ongoing program. Keck spectra of the host of candidate-1 in MS1054 ($z = 0.83$) show it is an early-type cluster galaxy devoid of any emission lines, arguing that the transient from 2002 (its appearance in the difference image is shown as an inset) was a near-nuclear SN-Ia in a cluster-galaxy. The other events shown have similarly been classified as cluster, background and foreground events (Table 1).




FIGURE 3. Examples of recently obtained Keck spectra. Top: a faint SN candidate, the spectrum shown contains the sum of the SN and host light, and final extraction awaits galaxy host light subtraction after the SN has faded. Bottom: a SN host galaxy, the derived redshift, $z = 0.582$, shows this was a background event.